

Getting Ready for your BRAVO pH Monitoring Test “BRAVO Prep”

Date:	Location:
Arrival Time:	Procedure Time:

Please read through the instructions at least one week before your procedure.

BE PREPARED!

If you have any questions, please call **207-773-7964**.

A responsible adult will need to drive you home. Taxis, Ubers, buses, medical ride services, or walking are not options unless a responsible adult accompanies you. You will not be able to drive for the rest of the day.

Plan on being at our facility for about 1 ½ hours from arrival to departure (although this may vary depending on the day/procedure/location – ex. approx. 2.5 hrs for MMC)

For **appointments with a scheduled arrival time of 4:00 pm or later**, have your ride plan to be at the facility **no later than 5:30 pm**.

If you are unable to keep your appointment, please call at least 3 business days prior to your procedure to reschedule. We reserve the right to charge a penalty fee for patients who “no show” or cancel without adequate notice.

NOTHING BY MOUTH 3 HOURS BEFORE YOUR PROCEDURE

Medication Instructions

- Aside from the medications below, we recommend you take all home medications as usual with a small amount of water, at least 3 hours before your arrival time.
- If you take blood thinners (Eliquis[®], Plavix[®], Warfarin, Xarelto[®], etc.) and haven't been given directions to stop them by our office or your prescribing physician's office, please call us to clarify. It is ok to continue your Aspirin.
- Take ½ your morning insulin dose. Bring the other ½ with you.

Stop the following medications until after your procedure as directed below:

7 days before	Weekly dosed GLP-1 agonists: Adipex-P [®] /Lomaira [®] (Phentermine), Bydureon [®] BCise [®] (Exenatide ER), Mounjaro [®] /Zepbound [®] (Tirzepatide), Ozempic [®] /Wegovy [®] (Semaglutide), Qsymia [®] (Phentermine/Topiramate), Trulicity [®] (Dulaglutide)
Day of procedure	Daily dosed GLP-1 agonists: Adlyxin [®] (Lixisenatide), Byetta [®] (Exenatide IR), Rybelsus [®] (Semaglutide), Saxenda [®] /Victoza [®] (Liraglutide) Morning dose of oral diabetic medication (Bring it with you)

NOTHING BY MOUTH 3 HOURS BEFORE YOUR PROCEDURE

Do you have questions? Contact our office at 207-773-7964.

Office is open 8:30 am – 4:30 pm Monday - Friday

Getting Ready

- ONE WEEK BEFORE** the test, stop the following medications:
 - **Aciphex, Nexium, Prevacid, Prilosec, and Protonix.**
- 24 HOURS BEFORE** the test, stop the following medications:
 - **Axid, Cimetidine, Pepcid, Tagamet, Zantac.**
 - 1 - 8.3 oz. container of **GaviLAX or MiraLAX®** powder
- You may take antacid (Tums) up 8 hours prior to your test.
- Call the office if you are pregnant, have an internal cardiac defibrillator or an artificial heart valve, had a recent cardiac procedure or a new cardiac health concern, you have an allergy to **nickel**.
- Review the Medication Instructions on the prior page.

Day of Your Procedure

8 Hours Prior to your Procedure

- No solid foods until after your procedure.
- You may drink clear liquids up until 3 hours before your procedure.

Examples of Clear Liquids*

Popsicles ~ Coconut water (not milk) ~ Gatorade® ~ Kool-Aid® Strained fruit juice
(apple or white grape) ~ Jell-O®
coffee and tea without dairy products/creamer ~ Soft Drinks

***No Red/Blue/Purple or items with sediment/solids**

Your prep is one of the most important parts of the procedure! Not following these steps may require rescheduling.

NOTHING BY MOUTH 3 HOURS BEFORE YOUR PROCEDURE

What are the risks of an endoscopy with BRAVO test?

Endoscopy is a general medical procedure. Although rare, all procedures have some risk.

1. The instrument may tear the wall of the throat, stomach or upper intestine, and may require surgery.
2. There may be adverse drug reactions or heart/breathing complications.
3. Bleeding (Bleeding may occur, usually when tissue or polyps are removed).
4. There may be discomfort during and after the procedure.

Do you have questions? Contact our office at 207-773-7964.

Office is open 8:30 am – 4:30 pm Monday - Friday

NOTHING BY MOUTH 3 HOURS BEFORE YOUR PROCEDURE